



Human Motor Neurons (iPSC-derived)

SKU: 40HU-005; 40HU-006

Product Description

Spinal motor neurons (MNs) are a highly specialized type of neurons that reside in the ventral horns and project axons to muscles to control their movement. Neurodegenerative diseases, such as spinal muscular atrophy (SMA), amyotrophic lateral sclerosis (ALS), Charcot-Marie-Tooth and poliomyelitis disease are a result of the progressive degeneration of motor neurons [1]. Furthermore, motor neurons derived from normal, or patient induced pluripotent stem cells (iPSCs) enable the generation of cell models with features relevant to human physiology, thus making it a valuable tool for biochemical analysis, disease modelling and other broad range of clinical applications [2,3].

Product Details

Catalog Number: 40HU-005; 40HU-006

Organism: Homo Sapiens, Human

Cell Type: Brain Cell

Tissue: N/A

Disease: Normal (40HU-005), ALS (40HU-006)

Package Size: 1 x 10⁶ cells/vial, 2 x 10⁶ cells/vial and 4 x 10⁶ cells/vial

Passage Number: P0

Growth Properties: Adherent

Associated Media: Motor Neuron Culture Medium Kit (MD-0022);

Motor Neuron Activity Medium Kit (MD-0118)

Storage Conditions & Shipment

Product Format/Shipped: Cryopreserved

Storage: Liquid Nitrogen

PRODUCT SHEET

For Research Use Only

iXCells Biotechnologies USA, Inc.

United States

www.ixcellsbiotech.com

Customer Support

E. supports@ixcellsbiotech.com

E. orders@ixcellsbiotech.com

P. +1 (858) 412 5988


RELATED PRODUCTS



Human Motor Neurons (iPSC-derived, TDP43 mutation,

M337V, HET)

SKU: 40HU-105-HET



Human Motor Neurons (iPSC-derived, TDP43 mutation, M337V, HET)

SKU: 40HU-105-HET

ı





Overview of Human Motor Neurons (iPSC-derived)

iXCells Biotechnologies is proud to provide the world's first fully differentiated and functional human iPSC-derived motor neurons that display typical neuronal morphology and express all key markers of motor neurons, e.g., HB9 (MNX1), ISL1, ChAT (Figure 1) when cultured in the Motor Neuron Culture Medium Kit (Cat# MD-0022-100ML). Moreover, whole cell patch clamp revealed that when cultured in Motor Neuron Activity Medium Kit (Cat# MD-0118-100ML) over 65% of the neurons exhibited mature spiking, and over 35% of the neurons had spontaneous activity at a holding potential of -60 mV (Figure 2) indicating the presence of a highly mature population of neurons.





Figure 1. A) Immunofluorescence staining showing HB9 and ChAT positive cells on day 2 and 7 in culture respectively. **B)** Flow cytometry measurements demonstrate >85% HB9 and >90% ISL1 positive cells on day 1-2.



-100

0.3

-100


-120

0

30

time(sec)

Figure 2. A) An image of the cells during the experiment. B) The Na/K currents were recorded in voltage clamp mode with test potentials of -100 mV to 90 mV. C) EPSCs were recorded in voltage clamp mode while clamping the cell at -60 mV. D) Evoked potentials were recorded in current clamp mode with 3pA current steps. E) Spontaneous activity was recorded in current clamp mode with a current injection needed for a membrane potential of -45 mV.



India Contact:
Life Technologies (India) Pvt. Ltd. 306, Aggarwal City Mall, Opposite M2K Pitampura, Delhi – 110034 (INDIA).
Ph: +91-11-42208000, 42208111, 42208222, Mobile: +91-9810521400, Fax: +91-11-42208444
Email: customerservice@lifetechindia.com Website: www.lifetechindia.com

0.6 0.7

time(sec)





Protocols

This procedure describes the steps required for thawing and plating hiPSC-Derived Motor Neurons using iXCells™ Motor Neuron Culture Medium Kit (Cat# MD-0022-100ML). The same procedure applies in the case of Motor Neuron Activity Medium Kit (Cat# MD-0118-100ML).

REQUIRED REAGENTS AND MATERIALS

IXCELLS™ MOTOR NEURON CULTURE MEDIUM KIT (CAT# MD-0022-100ML) INCLUDES:

		VOLUME
MOTOR NEURON CULTURE MEDIUM A	MD-0022A-100ML	100 ML
MOTOR NEURON CULTURE MEDIUM B	MD-0022B-20UL	20 µL

The motor neuron culture medium A and B can be stored at 2-8°C for 1 month and at -20°C for 12 months.

IXCELLS™ MOTOR NEURON ACTIVITY MEDIUM KIT (CAT# MD-0118-100ML) INCLUDES:

		VOLUME
MOTOR NEURON ACTIVITY MEDIUM A	MD-0118A-100ML	100 ML
MOTOR NEURON ACTIVITY MEDIUM B	MD-0118B-20UL	20 µL

The motor neuron activity medium A and B can be stored at 2-8°C for 1 month and at -20°C for 12 months.

Mono-culture of hiPSC-Derived Motor Neurons

The following protocol is based on 48-well plate format

 Prepare the coating vessel before thawing the cells. Please refer to the iXCell's <u>coating</u> protocol.

Note: Upon receipt of the frozen cells, it is recommended to thaw the cells and initiate the culture immediately in order to retain the highest cell viability.

- 2. To thaw the cells, put the vial in 37°C water bath with gentle agitation for ~1-2 minutes. Keep the cap out of water to minimize the risk of contamination.
- Pipette the cells into a 15 mL conical tube with 5 mL Motor Neuron Culture Medium A (Cat# MD-0022A-100ML).
- 4. Centrifuge at 600g for 5 minutes at room temperature. In the meantime, prepare a mix of Motor Neuron Culture Medium A with Motor Neuron Culture Medium B (Cat# MD-0022B-20UL; 1 in 1000) for plating. e.g. for 5 mL Motor Neuron Culture Medium A add 5 μL Motor Neuron Culture Medium B.

Note: Please spin down the vial of Motor Neuron Culture Medium B before opening it.





- 5. Remove the supernatant and re-suspend the cells in previously made mix of Motor Neuron Culture Medium A and B.
- 6. Seed the cells on precoated plate at the desired density. Incubate in 37°C CO₂ incubator overnight. Perform **complete media change** to remove the Motor Neuron Culture Medium B the next day.

Note: We recommend using a pipette when performing complete media change instead of aspirating.

Note: We recommend seeding 100-300K cells per cm² depending on the application. Cell debris may be observed after cell recovery because the cryopreserved neurons are fragile. Refer to the CoA of each lot to determine the seeding density for your experiment.

7. Perform **half medium** change every 2-3 days with Motor Neuron Culture Medium A only. Most of the cells should express high levels of HB9 and ISL1 1-2 days after thaw, and express high levels of ChAT and MAP2 7-10 days after thaw.

Note: Pure motor neurons tend to aggregate and detach from the plates. Change 50% of the medium with extra care to avoid cell loss.

Co-culture of hiPSC-Derived Motor Neurons with Astrocytes

The following protocol is based on 48-well plate format.

- Thaw a vial of iPSC-derived motor neuron as described previously, in Motor Neuron Culture Medium A (Cat# MD-0022A-100ML) with Motor Neuron Culture Medium B (Cat# MD-0022B-20UL; 1 in 1000). Seed the cells on precoated plate at the desired density. Incubate in 37°C CO₂ incubator overnight.
- Perform <u>complete media change</u> to remove the Motor Neuron Culture Medium B the next day.

Note: We recommend using a pipette when performing complete media change instead of aspirating.





- 3. Thaw a vial of Astrocyte (Cat# 40HU-008, check iPSC-derived Astrocyte product page for recovery details) and seed the cells on top of the motor neurons at 2x10⁴ cells per well (48-well plate format), in Motor Neuron Culture Medium A with 1% FBS. We recommend adding astrocytes 1 to 7 days post plating the iPSC derived motor neurons.
- 4. Incubate in 37°C CO₂ incubator overnight.
- **5.** Perform half medium change every 2-3 days (keeping the 1% FBS to support the astrocytes).





References

[1] Brady ST. (1993). "Motor neurons and neurofilaments in sickness and in health. Cell. 9;73(1):1-3.

[2] Dolmetsch R, Geschwind DH. (2011) "The human brain in a dish: the promise of iPSC-derived neurons". Cell. 145(6):831-4.

[3] Payne NL, Sylvain A, O'Brien C, Herszfeld D, Sun G, Bernard CC. (2015) "Application of human induced pluripotent stem cells for modeling and treating neurodegenerative diseases." Nat Biotechnology. 25;32(1):212-28.

Disclaimers

This product is for laboratory research purposes only and should not be used in humans. While iXCells Biotechnologies strives to provide accurate and current information on this product sheet, we do not guarantee its accuracy. Citations from scientific literature and patents are for informational purposes and may not be confirmed as accurate.

You are responsible for the safe storage, handling, and usage of this product, and iXCells Biotechnologies is not liable for any damages or injuries resulting from its receipt or use. While we make reasonable efforts to ensure the authenticity and reliability of deposited strains, we are not liable for damages arising from misidentification or misrepresentation of cultures.

Copyright

© iXCells Biotechnologies USA, Inc. 2023. All rights reserved.

Contact Information & Customer Support

iXCells Biotechnologies USA, Inc.

United States

www.ixcellsbiotech.com

Technical Supports. supports@ixcellsbiotech.com

Product Sales. orders@ixcellsbiotech.com or contact your local distributor

US Phone. +1 (858) 412 5988

Email: customerservice@lifetechindia.com Website: www.lifetechindia.com