



Overview

beta-Defensin 2 (BD-2), Human; **Synonyms**

> Defensins (alpha and beta) are cationic peptides with a broad spectrum of antimicrobial activity that comprise an important arm of the innate immune system. The a-defensins are distinguished from the ß-defensins by the pairing of their three disulfide bonds. To date, four human ß-defensins have been identified; BD-1, BD-2, BD-3 and BD-4. ß-defensins are expressed on some leukocytes and at epithelial surfaces. In addition to their direct antimicrobial activities, they are chemoattractant towards immature dendritic cells and

> memory T cells. The ß-defensin proteins are expressed as the C-terminal portion of precursors and are released by proteolytic cleavage of a signal sequence and, in the case of BD-1 (36 a.a.), a propeptide region. ß-defensins contain a six-cysteine motif that forms three intra-molecular disulfide bonds. ß-Defensins are 3-5 kDa peptides ranging in size

from 33-47 amino acid residues.

Species Human E. coli Source

Description

Fully biologically active when compared to standard. The biological activity determined by **Biological Activity** a chemotaxis bioassay using immature human dendritic cells is in a concentration range

of 10-100 ng/ml.

GIGDPVTCLK SGAICHPVFC PRRYKQIGTC GLPGTKCCKK Sequence

Properties

Measured Molecular Approximately 4.3 kDa, a single non-glycosylated polypeptide chain containing 41 amino

Weight acids.

Purity > 98 % by SDS-PAGE and HPLC analyses.

Lyophilized from a 0.2 µm filtered concentrated solution in 20 mM PBS, pH 7.4, 130 mM **Formulation**

NaCl.

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at d -20 °C. Further dilutions should be made in appropriate buffered

solutions.

Endotoxin Level Less than 1 EU/µg of rHuBD-2 as determined by LAL method.

Physical Appearance

Storage

Reconstitution

Sterile Filtered White lyophilized (freeze-dried) powder.

This material is for research, laboratory or further evaluation purposes. NOT FOR Usage

HUMAN USE.

This lyophilized preparation is stable at 2-8 °C, but should be kept at -20 °C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one

week at 2-8 °C. For maximal stability, apportion the reconstituted preparation into working

aliquots and store at -20 °C to -70 °C. Avoid repeated freeze/thaw cycles.

India Contact:

Life Technologies (India) Pvt. Ltd.

306, Aggarwal City Mall, Opposite M2K Pitampura,

Delhi - 110034 (INDIA).

Mobile: +91-9810521400, Ph: +91-11-42208000 Email: customerservice@lifetechindia.com

Web: www.lifetechindia.com