



Overview

Synonyms MCP-3/CCL7, Human;

MCP2 and CCL7 are two monocyte chemotactic proteins produced by human MG63 osteosarcoma cells. Both MCP2 and CCL7 are members of the CC family of chemokines and share 62% and 71% amino acid sequence identity, respectively, with MCP1. CCL7 also shares 58% amino acid identity with MCP2. Similarly to other CC chemokines, all

three MCP proteins are monocyte chemoattractants. In addition, the three MCPs can chemoattract activated NK cells as well as CD4+ and CD8+ T lymphocytes. All three cytokines have also been shown to attract eosinophils and induce histamine secretion

from basophils.

Species Human Source E. coli

Description

Fully biologically active when compared to standard. The biologically active determined by

Biological Activity a chemotaxis bioassay using human monocytes is in a concentration range of 10-100

ng/ml.

Sequence QPVGINTSTT CCYRFINKKI PKQRLESYRR TTSSHCPREA VIFKTKLDKE ICADPTOKWV ODFMKHLDKK TOTPKL

Properties

Measured Molecular Approximately 9.0 kDa, a single, non-glycosylated polypeptide chain containing 76 amino

Weight acids.

Purity > 97 % by SDS-PAGE and HPLC analyses.

Formulation Lyophilized from a 0.2 μm filtered concentrated solution in 20 mM PB, pH 7.4, 150 mM

NaCI.

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working

aliquots and stored at d -20 °C. Further dilutions should be made in appropriate buffered

solutions.

Endotoxin Level

Reconstitution

Less than 1 EU/μg of rHuMCP-3/CCL7 as determined by LAL method.

Physical Appearance

Storage

Sterile Filtered White lyophilized (freeze-dried) powder.

Usage This material is for research, laboratory or further evaluation purposes. NOT FOR

HUMAN USE.

This lyophilized preparation is stable at 2-8 °C, but should be kept at -20 °C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one

week at 2-8 °C. For maximal stability, apportion the reconstituted preparation into working

aliquots and store at -20 °C. Avoid repeated freeze/thaw cycles.

India Contact:

Life Technologies (India) Pvt. Ltd.

306, Aggarwal City Mall, Opposite M2K Pitampura,

Delhi - 110034 (INDIA).

Mobile: +91-9810521400, Ph: +91-11-42208000 Email: customerservice@lifetechindia.com

Web: www.lifetechindia.com