



Overview

Synonyms LIX/CXCL5 (93a.a.), Mouse;

The mouse homolog of ENA-78 is called LIX. ENA-78/LIX is a CXC chemokine that signals through the CXCR2 receptor. It is expressed in monocytes, platelets, endothelial cells, and mast cells. ENA-78/LIX is a chemoattractant for neutrophils. The three naturally occurring variants of human ENA-78; ENA 5-78, ENA 9-78 and ENA 10-78, contain 74,

70, and 69 amino acid residues, respectively, and possess the same biological activity. ENA-78/LIX contains the four conserved cysteine residues present in CXC chemokines,

and also contains the 'ELR' motif common to CXC chemokine that bind to the CXCR1 and

CXCR2 receptors.

Species Mouse Source E. coli

Description

Fully biologically active when compared to standard. The biological activity determined by

Biological Activity a chemotaxis bioassay using human peripheral blood neutrophils is in a concentration of

10-100 ng/ml.

APSSVIAATE LRCVCLTVTP KINPKLIANL EVIPAGPQCP

Sequence TVEVIAKLKN QKEVCLDPEA PVIKKIIQKI LGSDKKKAKR NALAVERTAS

VQ

Properties

Measured Molecular Approximately 9.8 kDa, a single, non-glycosylated polypeptide chain containing 92 amino

Weight acids.

Purity > 97 % by SDS-PAGE and HPLC analyses.

Formulation Lyophilized from a 0.2 μm filtered concentrated solution in 20 mM PB, pH 7.4, 150 mM

NaCI.

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA

Reconstitution to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working

aliquots and stored at d -20 °C. Further dilutions should be made in appropriate buffered

solutions.

Endotoxin Level Less than 1 EU/µg of rMuLIX/CXCL5 as determined by LAL method.

Physical Appearance

Storage

Sterile Filtered White lyophilized (freeze-dried) powder.

Usage This material is for research, laboratory or further evaluation purposes. NOT FOR

HUMAN USE.

This lyophilized preparation is stable at 2-8 °C, but should be kept at -20 °C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one

week at 2-8 °C. For maximal stability, apportion the reconstituted preparation into working

aliquots and store at -20 °C to -70 °C. Avoid repeated freeze/thaw cycles.

India Contact:

Life Technologies (India) Pvt. Ltd.

306, Aggarwal City Mall, Opposite M2K Pitampura,

Delhi - 110034 (INDIA).

Mobile: +91-9810521400, Ph: +91-11-42208000 Email: customerservice@lifetechindia.com

Web: www.lifetechindia.com