

Overview

Synonyms	EDF, BCDFII, TRF
Description	<p>Interleukin-5 (IL-5), produced by mast cells, T cells and eosinophils, is responsible for the activities attributed to eosinophil differentiating factor, B cell growth factor II and T cell-replacing factor (TRF). It can increase production and mobilization of eosinophils and CD34+ progenitors from the bone marrow. IL-5 plays an important role in inducing cell-mediated immunity against parasitic infections and certain tumors. IL-5 also promotes differentiation of basophils and primes them for histamine and leukotriene release.</p> <p>Recombinant Murine IL-5 is homodimeric protein with molecular weight ranging from 25 to 40 kDa due to glycosylation.</p>
Species	Mouse
Source	CHO
Biological Activity	ED ₅₀ < 0.5 ng/ml, measured in a cell proliferation assay using TF-1 cells, corresponding to a specific activity of > 2×10 ⁶ units/mg.
Sequence	<p>MEIPMSTVVK ETLTQLSAHR ALLTSNETMR LPVPTHKNHQ</p> <p>LCIGEIFQGL DILKNQTVRG GTVEMLFQNL SLIKKYIDRQ</p> <p>KEKCGEERRR TRQFLDYLQE FLGVMSTEWA MEG</p>

Properties

Measured Molecular Weight	25-40 kDa, observed by non-reducing SDS-PAGE.
Purity	> 95% as analyzed by SDS-PAGE and HPLC.
Formulation	Lyophilized after extensive dialysis against PBS.
Reconstitution	Reconstituted in ddH ₂ O or PBS at 100 µg/ml.
Endotoxin Level	< 0.2 EU/µg, determined by LAL method.
Storage	Lyophilized recombinant Murine Interleukin 5 (IL-5) remains stable up to 6 months at lower than -70°C from date of receipt. Upon reconstitution, rmlIL-5 should be stable up to 1 week at 4°C or up to 2 months at -20°C.
Note	For research use only

India Contact:

Life Technologies (India) Pvt. Ltd.

306, Aggarwal City Mall, Opposite M2K Pitampura,
Delhi – 110034 (INDIA).

Mobile: +91-9810521400, Ph: +91-11-42208000

Email: customerservice@lifetechindia.com

Web: www.lifetechindia.com