

## Overview

<b>Synonyms</b>	BCGF, BCDF, B cell stimulating factor (BSF-1)
<b>Description</b>	<b>Interleukin-4 (IL-4)</b> is a pleiotropic cytokine regulates diverse T and B cell responses including cell proliferation, survival, and gene expression. It has important effects on the growth and differentiation of different immunologically competent cells. Interleukin-4 is produced by mast cells, T cells, and bone marrow stromal cells <sup>[1]</sup> . IL-4 regulates the differentiation of native CD4+ T cells (Th0 cells) into helper Th2 cells, and regulates the immunoglobulin class switching to the IgG1 and IgE isotypes. IL-4 has numerous important biological functions including stimulating B-cells activation, T-cell proliferation and CD4+ T-cells differentiation to Th2 cells. It is a key regulator in hormone control and adaptive immunity <sup>[2]</sup> . IL-4 also plays a major role in inflammation response and wound repair via activation of macrophage into M2 cells <sup>[3]</sup> . IL-4 is stabilized by three disulphide bonds forming a compact globular protein structure <sup>[4]</sup> . Four alpha-helix bundle with left-handed twist is dominated half of the protein structure with 2 overhand connections and fall into a 2-stranded anti-parallel beta sheet <sup>[5]</sup> .
<b>Species</b>	Mouse
<b>Source</b>	CHO
<b>Biological Activity</b>	ED <sub>50</sub> < 2 ng/ml, measured in a cell proliferation assay using murine HT-2 cells, corresponding to a specific activity of >5 x 10 <sup>5</sup> units/mg.
<b>Sequence</b>	HIHGCDKNHL REIIGILNEV TGEETPCTEM DVPNVLTATK NTTESELVCR ASKVLRIFYI KHGKTPCLKK NSSVLMELQR LFRAFRCLDS SISCTMNESK STSLKDFLES LKSIMQMDYS

## Properties

<b>Measured Molecular Weight</b>	15 kDa, observed by reducing SDS-PAGE.
<b>Purity</b>	> 95% as analyzed by SDS-PAGE and HPLC.
<b>Formulation</b>	Lyophilized after extensive dialysis against PBS.
<b>Reconstitution</b>	Reconstituted in ddH <sub>2</sub> O or PBS at 100 µg/ml.
<b>Endotoxin Level</b>	< 0.2 EU/µg, determined by LAL method.
<b>Storage</b>	Lyophilized recombinant <b>Murine Interleukin 4 (IL-4)</b> remains stable up to 6 months at lower than -70°C from date of receipt. Upon reconstitution, rmlL-4 should be stable up to 1 week at 4°C or up to 2 months at -20°C.
<b>Note</b>	For research use only

India Contact:

**Life Technologies (India) Pvt. Ltd.**

306, Aggarwal City Mall, Opposite M2K Pitampura,  
Delhi – 110034 (INDIA).

Mobile: +91-9810521400, Ph: +91-11-42208000

Email: [customerservice@lifetechindia.com](mailto:customerservice@lifetechindia.com)

Web: [www.lifetechindia.com](http://www.lifetechindia.com)