

Overview

Synonyms	CXCL1, GRO \pm , NAP-3, GRO1, KC (murine), CINC (rat)
Description	GRO/MGSA/CXCL1 has chemotactic activity for neutrophils. It may play a role in inflammation and exerts its effects on endothelial cells in an autocrine fashion. All three isoforms of GRO are CXC chemokines that can signal through the CXCR1 or CXCR2 receptors. GRO expression is inducible by serum or PDGF and/or by a variety of inflammatory mediators, such as IL-1 and TNF, in monocytes, fibroblasts, melanocytes and epithelial cells. In certain tumor cell lines, GRO is expressed constitutively.
Accession No	P14095
Species	Rat
Source	HEK 293
Biological Activity	Active, measured in a functional assay using HUVEC cells.
Sequence	APVANELRCQ CLQTVAGIHF KNIQSLKVMP PGPHTQTEV IATLKNGREA CLDPEAPMVQ KIVQKMLKGV PK

Properties

Measured Molecular Weight	~7.8 kDa, observed by reducing SDS-PAGE.
Purity	> 95% as analyzed by SDS-PAGE.
Formulation	Lyophilized after extensive dialysis against PBS.
Reconstitution	Reconstituted in ddH ₂ O or PBS at 100 μ g/ml.
Endotoxin Level	< 0.2 EU/ μ g, determined by LAL method.
Storage	Lyophilized recombinant Rat GRO/MGSA/CXCL1 remains stable up to 6 months at lower than -70°C from date of receipt. Upon reconstitution, Rat GRO/MGSA/CXCL1 should be stable up to 1 week at 4°C or up to 2 months at -20°C.
Note	For research use only

India Contact:

Life Technologies (India) Pvt. Ltd.

306, Aggarwal City Mall, Opposite M2K Pitampura,
Delhi – 110034 (INDIA).

Mobile: +91-9810521400, Ph: +91-11-42208000

Email: customerservice@lifetechindia.com

Web: www.lifetechindia.com