

nTaq-Tenuto

Cat.#	Size	Size
P225A	250 units	5 units/μl
P250A	500 units	5 units/μl

Store at -20°C

Supplied with: 10X nTaq-Tenuto Buffer (Mg²⁺ plus)
dNTP Mixture (2 mM each)
GC Melt I
GC Melt II
Sterile water

India Contact:

Life Technologies (India) Pvt. Ltd.
306, Aggarwal City Mall, Opposite M2K Pitampura,
Delhi – 110034 (INDIA).
Ph: +91-11-42208000, 42208111, 42208222
Mobile: +91-9810521400
Fax: +91-11-42208444
Email: customerservice@lifetechindia.com
Web: www.lifetechindia.com

Product description

nTaq-Tenuto is nTaq (Cat.# P025, P050) supplemented with 3'→5' proofreading activity and a PCR enhancing factor for improved efficiency and fidelity. nTaq-Tenuto DNA Polymerase can be used to amplify DNA longer than 10 kb, which is difficult with common Taq polymerases alone. Thus, this product is improved in both fidelity (> 2 fold) of PCR products and amplification efficiency of longer PCR products.

Characteristics

- Molecular weight: 94 kDa
- Error rate: 3.0 X 10⁻⁶
- Thermal stability: Half life of 40 min at 95°C
- A-tail formation at 3' ends of amplified DNA products.

Applications

- Amplification of long DNA fragments (>5~15 kb)
- Amplification of high-complexity template DNA such as cDNA and genomic DNA
- Primer extension
- Colony PCR
- Multiplex PCR
- Labeling of DNA fragments with radioactive-isotopes
- Nucleotide sequencing

For Research Use Only. Not for use in diagnostic procedures.

ISO9001 ISO14001 ISO13485

Quality control

- Purity: >99% on SDS-PAGE
- Endonuclease-free
- Exonuclease-free
- RNase free
- Inhibitor-free

Unit definition

One unit is defined as the amount of enzyme required to incorporate 10 nmol of dNTP into acid insoluble materials in 30 min at 74°C in a 50-μl reaction mixture (20 mM Tris-HCl/pH 8.8, 50 mM KCl, 2.5 mM MgCl₂, 10 mM β-mercaptoethanol, 12.5 μg of calf thymus DNA).

Storage buffer

20 mM Tris-HCl (pH 7.9), 100 mM KCl, 0.1 mM EDTA, 1 mM DTT, 0.5% NP-40, 0.5% Tween-20, 50% glycerol.

10X nTaq-Tenuto Buffer (Mg²⁺ plus)

Mg²⁺ plus buffer: Containing 20 mM Mg²⁺

nTaq-Tenuto

Cat.#	Size	Size
P225A	250 units	5 units/μl
P250A	500 units	5 units/μl

Store at -20°C

Supplied with: 10X nTaq-Tenuto Buffer (Mg²⁺ plus)
dNTP Mixture (2 mM each)
GC Melt I
GC Melt II
Sterile water

Product description

nTaq-Tenuto is nTaq (Cat.# P025, P050) supplemented with 3'→5' proofreading activity and a PCR enhancing factor for improved efficiency and fidelity. nTaq-Tenuto DNA Polymerase can be used to amplify DNA longer than 10 kb, which is difficult with common Taq polymerases alone. Thus, this product is improved in both fidelity (> 2 fold) of PCR products and amplification efficiency of longer PCR products.

Characteristics

- Molecular weight: 94 kDa
- Error rate: 3.0 X 10⁻⁶
- Thermal stability: Half life of 40 min at 95°C
- A-tail formation at 3' ends of amplified DNA products.

Applications

- Amplification of long DNA fragments (>5~15 kb)
- Amplification of high-complexity template DNA such as cDNA and genomic DNA
- Primer extension
- Colony PCR
- Multiplex PCR
- Labeling of DNA fragments with radioactive-isotopes
- Nucleotide sequencing

For Research Use Only. Not for use in diagnostic procedures.

ISO9001 ISO14001 ISO13485

Quality control

- Purity: >99% on SDS-PAGE
- Endonuclease-free
- Exonuclease-free
- RNase free
- Inhibitor-free

Unit definition

One unit is defined as the amount of enzyme required to incorporate 10 nmol of dNTP into acid insoluble materials in 30 min at 74°C in a 50-μl reaction mixture (20 mM Tris-HCl/pH 8.8, 50 mM KCl, 2.5 mM MgCl₂, 10 mM β-mercaptoethanol, 12.5 μg of calf thymus DNA).

Storage buffer

20 mM Tris-HCl (pH 7.9), 100 mM KCl, 0.1 mM EDTA, 1 mM DTT, 0.5% NP-40, 0.5% Tween-20, 50% glycerol.

10X nTaq-Tenuto Buffer (Mg²⁺ plus)

Mg²⁺ plus buffer: Containing 20 mM Mg²⁺

India Contact:

Life Technologies (India) Pvt. Ltd.
306, Aggarwal City Mall, Opposite M2K Pitampura,
Delhi – 110034 (INDIA).
Ph: +91-11-42208000, 42208111, 42208222
Mobile: +91-9810521400
Fax: +91-11-42208444
Email: customerservice@lifetechindia.com
Web: www.lifetechindia.com

Your Molecular & Cell Technology Partner

Standard PCR conditions

- PCR mixture^a

10X nTaq-Tenuto Buffer (Mg ²⁺ plus)	2 μl
nTaq-Tenuto DNA Polymerase ^b (5 units/μl)	0.2 μl
dNTP mixture (2 mM each, final conc., 200 μM each)	2 μl
Template DNA ^c (0.1~500 ng/μl)	1 μl
Primer 1 (5 pmole/μl)	1 μl
Primer 2 (5 pmole/μl)	1 μl
Sterile water	up to 20 μl
^a Assemble the reaction mixture on ice	
^b Add the PCR polymerase at the final step	
^c Plasmid DNA, 0.1 ng~30 ng; genomic DNA, 50 ng~500 ng	

- PCR cycle

Initial denaturation	95°C	2 min
Denaturation	95°C	30 sec
Annealing ^a	55°C~65°C	30~60 sec
Elongation	72°C	1 min/kb
Number of cycles	25~35 times	
Final elongation	72°C	5 min
When cycles are over, keep the reaction mixture at 4°C; may add 10 mM EDTA until use to prevent DNA degradation.		
^a Recommended annealing temperatures is 5 to 10°C below the lower T _m of the two primers used.		

Your Molecular & Cell Technology Partner

Standard PCR conditions

- PCR mixture^a

10X nTaq-Tenuto Buffer (Mg ²⁺ plus)	2 μl
nTaq-Tenuto DNA Polymerase ^b (5 units/μl)	0.2 μl
dNTP mixture (2 mM each, final conc., 200 μM each)	2 μl
Template DNA ^c (0.1~500 ng/μl)	1 μl
Primer 1 (5 pmole/μl)	1 μl
Primer 2 (5 pmole/μl)	1 μl
Sterile water	up to 20 μl
^a Assemble the reaction mixture on ice	
^b Add the PCR polymerase at the final step	
^c Plasmid DNA, 0.1 ng~30 ng; genomic DNA, 50 ng~500 ng	

- PCR cycle

Initial denaturation	95°C	2 min
Denaturation	95°C	30 sec
Annealing ^a	55°C~65°C	30~60 sec
Elongation	72°C	1 min/kb
Number of cycles	25~35 times	
Final elongation	72°C	5 min
When cycles are over, keep the reaction mixture at 4°C; may add 10 mM EDTA until use to prevent DNA degradation.		
^a Recommended annealing temperatures is 5 to 10°C below the lower T _m of the two primers used.		