



Product Information Sheet

M5865 PhytoReady™ Murashige & **Skoog Germination/Initiation** Media

Properties:

Form: Gel

Appearance:

Plant Tissue Culture Media Application:

Solubility: Water

Typical Working Concentration:

22.44q/L

Storage Temp:

2-8 °C

Other Notes:

Contains the macro- and micronutrients and vitamins as described by Murashige and

Skoog (1962). Also contains 10 g/L Sucrose and 8 g/L Agar. Premixed, sterilized, and

dispensed into 25x150 mm polycarbonate culture tubes.

Formula [mg/L]:

Ammonium Nitrate	1650
Boric Acid	6.2
Cupric Sulfate•5H ₂ O	0.025
Ferrous Sulfate	27.8
Magnesium Sulfate, Anhydrous	180.7
Manganese Sulfate•H₂O	16.9
Molybdic Acid (Sodium Salt)•2H ₂ O	0.25
Potassium Iodide	0.83
Potassium Nitrate	1900

Potassium Phosphate Monobasic	170
Zinc Sulfate•7H₂O	8.6
myo-Inositol	100
Nicotinic Acid (Free Acid)	1
Pyridoxine•HCl	1
Sucrose	10,000
Thiamine•HCI	10
Agar, Micropropagation Grade	8000

Application Notes:

PhytoReady™ Murashige & Skoog Multiplication Media is a ready-to-use plant tissue culture medium gelled in polycarbonate culture tubes (product C2035). Murashige and Skoog (MS)-media used in this product was developed to culture tobacco, and in general works well supporting the growth of herbaceous (non-woody) plant species. It is the most commonly used media in plant tissue culture. This medium contains sucrose as a carbohydrate source, and agar as a gelling agent. Each tube contains 20 mL of media. After 30-60 days, tissue can be subcultured to multiply the tissue into more shoots, or it can be applied to pre-transplant or rooting medium to encourage root organogenesis and proliferation on tissue. It can take weeks to many months for plant tissue to first become acclimated to growth in tissue culture with plant growth regulators. It can be erratic with very substantial growth and slow growth. Therefore, there are no plant growth regulators in this medium. Unwrap the tubes, and aseptically add disinfected*** tissue to the medium, and store at 25°C under sufficient lighting at 16 hours of light/8 hours of dark intervals. Young juvenile tissue is best to start from.

Please Note: While PhytoTechnology Laboratories® tests each lot of this product with two or more plant cell/ tissue culture lines, it is the sole responsibility of the purchaser to determine the appropriateness of this product for the specific plants that are being cultured and applications that are being used. ***Tissue is generally disinfected for 5-15 minutes in a solution of bleach diluted anywhere from 1:10 to 1:2 with a few drops of Tween 20 (P720) added per 50 mL's and gently mixed. It should then be rinsed with 3-4 volumes of autoclaved/pressure cooked sterile water. For more information on explant decontamination, please look at Prod. No. E2620 and the associated manual. These are merely guidelines, and your tissue may require more or less time in more or less concentrated bleach solutions. The goal of tissue disinfection is to find the least harsh solution and amount of time in that solution

PhytoTech Labs Inc.

14610 W 106th St. Lenexa, KS 66215





Product Information Sheet

that will remove fungi and bacteria, yet not induce plant cell and tissue death. Browning of the tissue can sometimes occur during disinfection. This can be reduced by reducing the time incubated in the bleach solution and the concentration of diluted bleach. The spread of browning on the tissue can also be reduced by dipping tissue in the Antioxidant solution (Product No. A126) following the final sterile water rinse.

References:

Murashige T and F Skoog. (1962) A revised medium for rapid growth and bioassays with tobacco tissue cultures. Physiol. Plant. 15:473-497

PhytoTech Labs Inc.